



## Short Communication

## Widespread occurrence of honey bee pathogens in solitary bees

Jorgen Ravoet<sup>a,\*</sup>, Lina De Smet<sup>a</sup>, Ivan Meeus<sup>b</sup>, Guy Smaghe<sup>b</sup>, Tom Wenseleers<sup>c</sup>, Dirk C. de Graaf<sup>a</sup><sup>a</sup> Laboratory of Zoophysiology, Ghent University, B-9000 Ghent, Belgium<sup>b</sup> Laboratory of Agrozoology, Ghent University, B-9000 Ghent, Belgium<sup>c</sup> Laboratory of Socioecology and Social Evolution, K.U. Leuven, B-3000 Leuven, Belgium

## ARTICLE INFO

## Article history:

Received 14 April 2014

Accepted 26 August 2014

Available online 6 September 2014

## Keywords:

Pathogens

Honey bees

Lake Sinai Virus

*Crithidia mellifica**Apis mellifera* Filamentous Virus

## ABSTRACT

Solitary bees and honey bees from a neighbouring apiary were screened for a broad set of putative pathogens including protists, fungi, spiroplasmas and viruses. Most sampled bees appeared to be infected with multiple parasites. Interestingly, viruses exclusively known from honey bees such as *Apis mellifera* Filamentous Virus and *Varroa destructor* Macula-like Virus were also discovered in solitary bees. A microsporidium found in *Andrena vaga* showed most resemblance to *Nosema thomsoni*. Our results suggest that bee hives represent a putative source of pathogens for other pollinators. Similarly, solitary bees may act as a reservoir of honey bee pathogens.

© 2014 Elsevier Inc. All rights reserved.

## 1. Introduction

There is a long tradition of studying the pathogens of the Western honey bee (*Apis mellifera*) and the list of honey bee pathogens has been expanded significantly from the sixties on. It became recently evident that common honey bee pathogens such as Deformed Wing Virus (DWV) can infect other bees as well (Furst et al., 2014; Levitt et al., 2013). Although several macroparasites of wild bees are well known (Westrich, 1990), reports on their microparasites are rather scarce. The few known solitary bee-specific parasites are fungi, including *Ascosphaera* spp. (Wynns et al., 2013) and *Antonospora scoticæ* (Fries et al., 1999).

The aim of this study was to investigate whether solitary bees sampled nearby an apiary harbour some of the known or recently discovered honey bee pathogens (Runckel et al., 2011). We screened for a broad set of parasitic micro-organisms, including fungi, protists, spiroplasmas and viruses.

## 2. Material and methods

## 2.1. Sample collection

For each species, three pooled samples of 10 bees were collected in 2012 at campus Sterre of Ghent University. *Osmia bicornis*, *Osmia cornuta* and *Heriades truncorum* were sampled at a bee hotel, close to an apiary. *Andrena vaga* and *Andrena ventralis* (only one

sample of 10 bees) were caught nearby their nest aggregations. These species were identified using suitable keys (Scheuchl, 1996; Schmid-Egger and Scheuchl, 1997). Three different honey bee colonies were simultaneously (within one week) sampled at the neighbouring apiary.

## 2.2. RNA and DNA extraction

Ten bees were homogenised in 5 ml PBS in the presence of glass beads. Total RNA was extracted from 100 µl supernatant using the RNeasy Lipid Tissue Kit (Qiagen). Using random hexamer primers, 1 µg RNA was retro-transcribed with the RevertAid First Strand cDNA Synthesis Kit (Thermo Scientific). DNA was extracted from 120 µl supernatant using the DNeasy Blood & Tissue Kit (Qiagen) according to the manufacturer's instructions for animal tissues.

## 2.3. PCR and BeeDoctor analysis

All PCR reaction mixtures contained: 2 µM of each primer (Table A1); 1.0 mM MgCl<sub>2</sub>; 0.2 mM dNTPs; 1.25 U Hotstar Taq DNA polymerase (Qiagen) and 1 µl cDNA or 3 µl DNA product (*Apis mellifera* Filamentous Virus (AmFV) and *Ascosphaera* spp. detection) using described PCR cycles (Table A1). For our developed primers we used: 94 °C for 15 min; [94 °C for 30 s, 50 °C (*Crithidia* spp. cytochrome *b*) or 55 °C (LSV Orf1) for 30 s, 72 °C for 1 min] 35 cycles, 72 °C for 10 min. Positive and negative controls were always included. Amplicons intended for Genbank submission were amplified with Hotstar High Fidelity Taq DNA polymerase

\* Corresponding author. Fax: +32 926 452 42.

E-mail address: [Jorgen.Ravoet@UGent.be](mailto:Jorgen.Ravoet@UGent.be) (J. Ravoet).

**Table 1**

Summary of pathogens detected in honey bees and solitary bees.

| Sample                | Date          | Virus                             | Crithidia spp.      | Neogregarinida  | Nosema spp.                        | Ascosphaera spp.        | Spiroplasma spp.     |
|-----------------------|---------------|-----------------------------------|---------------------|-----------------|------------------------------------|-------------------------|----------------------|
| Hive 3–1              | 4 April 2012  | ALPV, BQCV, LSV, AmFV             | –                   | <i>A. bombi</i> | <i>N. apis</i> , <i>N. ceranae</i> | –                       | –                    |
| Hive 5–1              | 4 April 2012  | ALPV, BQCV, LSV, AmFV             | <i>C. mellifica</i> | <i>A. bombi</i> | <i>N. apis</i> , <i>N. ceranae</i> | –                       | –                    |
| Hive 10–1             | 4 April 2012  | ALPV, BQCV, LSV, AmFV             | <i>C. mellifica</i> | <i>A. bombi</i> | <i>N. ceranae</i>                  | –                       | –                    |
| <i>O. cornuta</i> 1   | 4 April 2012  | BQCV, LSV, AmFV, VdMLV            | –                   | <i>A. bombi</i> | <i>N. ceranae</i>                  | +                       | –                    |
| Hive 3–2              | 16 April 2012 | ALPV, BQCV, LSV, AmFV             | <i>C. mellifica</i> | <i>A. bombi</i> | <i>N. apis</i> , <i>N. ceranae</i> | <i>A. apis</i>          | –                    |
| Hive 5–2              | 16 April 2012 | ALPV, BQCV, LSV, AmFV             | <i>C. mellifica</i> | <i>A. bombi</i> | <i>N. ceranae</i>                  | <i>A. apis</i>          | –                    |
| Hive 10–2             | 16 April 2012 | ALPV, BQCV, LSV, AmFV             | <i>C. mellifica</i> | <i>A. bombi</i> | <i>N. ceranae</i>                  | <i>A. apis</i>          | –                    |
| <i>O. cornuta</i> 2   | 16 April 2012 | BQCV, LSV, AmFV, VdMLV            | –                   | <i>A. bombi</i> | <i>N. ceranae</i>                  | <i>Ascosphaera</i> spp. | –                    |
| Hive 3–3              | 23 April 2012 | ALPV, BQCV, LSV, AmFV             | –                   | <i>A. bombi</i> | <i>N. apis</i> , <i>N. ceranae</i> | <i>A. apis</i>          | –                    |
| Hive 5–3              | 23 April 2012 | ALPV, BQCV, LSV, AmFV             | <i>C. mellifica</i> | <i>A. bombi</i> | <i>N. ceranae</i>                  | <i>A. apis</i>          | –                    |
| Hive 10–3             | 23 April 2012 | ALPV, BQCV, LSV, AmFV             | <i>C. mellifica</i> | <i>A. bombi</i> | <i>N. ceranae</i>                  | <i>A. apis</i>          | –                    |
| <i>O. cornuta</i> 3   | 24 April 2012 | BQCV, LSV, AmFV, VdMLV            | –                   | <i>A. bombi</i> | <i>N. ceranae</i>                  | <i>Ascosphaera</i> spp. | –                    |
| <i>A. vaga</i> 1      | 24 April 2012 | BQCV, LSV, AmFV                   | <i>C. bombi</i>     | <i>A. bombi</i> | <i>N. thomsoni</i>                 | –                       | –                    |
| <i>A. ventralis</i>   | 24 April 2012 | LSV, AmFV                         | –                   | <i>A. bombi</i> | <i>N. ceranae</i>                  | –                       | –                    |
| Hive 3–4              | 15 May 2012   | ALPV, BQCV, LSV, AmFV             | –                   | <i>A. bombi</i> | <i>N. apis</i> , <i>N. ceranae</i> | <i>A. apis</i>          | –                    |
| Hive 5–4              | 15 May 2012   | ALPV, BQCV, LSV, AmFV             | –                   | <i>A. bombi</i> | <i>N. ceranae</i>                  | –                       | –                    |
| Hive 10–4             | 15 May 2012   | ALPV, BQCV, LSV, AmFV             | <i>C. mellifica</i> | <i>A. bombi</i> | <i>N. ceranae</i>                  | <i>A. apis</i>          | –                    |
| <i>A. vaga</i> 2      | 8 May 2012    | LSV, SBV, AmFV                    | <i>C. bombi</i>     | <i>A. bombi</i> | <i>N. thomsoni</i>                 | –                       | –                    |
| <i>A. vaga</i> 3      | 8 May 2012    | LSV, AmFV                         | <i>C. bombi</i>     | <i>A. bombi</i> | <i>N. thomsoni</i>                 | –                       | –                    |
| Hive 3–5              | 31 May 2012   | ALPV, BQCV, DWV, LSV, VdMLV, AmFV | <i>C. mellifica</i> | <i>A. bombi</i> | <i>N. ceranae</i>                  | –                       | <i>S. melliferum</i> |
| Hive 5–5              | 31 May 2012   | ALPV, BQCV, DWV, LSV, VdMLV, AmFV | <i>C. mellifica</i> | <i>A. bombi</i> | <i>N. ceranae</i>                  | –                       | –                    |
| Hive 10–5             | 31 May 2012   | BQCV, DWV, LSV, VdMLV, AmFV       | <i>C. mellifica</i> | <i>A. bombi</i> | <i>N. ceranae</i>                  | –                       | –                    |
| <i>O. bicornis</i> 1  | 30 May 2012   | DWV, LSV, AmFV, VdMLV             | <i>C. bombi</i>     | <i>A. bombi</i> | <i>N. ceranae</i>                  | <i>Ascosphaera</i> spp. | <i>S. melliferum</i> |
| <i>O. bicornis</i> 2  | 30 May 2012   | DWV, LSV, AmFV, VdMLV             | <i>C. bombi</i>     | <i>A. bombi</i> | <i>N. ceranae</i>                  | <i>Ascosphaera</i> spp. | <i>S. apis</i>       |
| <i>O. bicornis</i> 3  | 30 May 2012   | DWV, LSV, AmFV, VdMLV             | <i>C. bombi</i>     | <i>A. bombi</i> | <i>N. ceranae</i>                  | <i>Ascosphaera</i> spp. | <i>S. melliferum</i> |
| Hive 3–6              | 9 July 2012   | ALPV, BQCV, DWV, LSV, VdMLV, AmFV | <i>C. mellifica</i> | <i>A. bombi</i> | <i>N. ceranae</i>                  | –                       | <i>S. melliferum</i> |
| Hive 5–6              | 9 July 2012   | DWV, LSV, VdMLV, AmFV             | <i>C. mellifica</i> | <i>A. bombi</i> | <i>N. ceranae</i>                  | <i>A. apis</i>          | <i>S. melliferum</i> |
| Hive 10–6             | 9 July 2012   | ALPV, BQCV, DWV, LSV, VdMLV, AmFV | <i>C. mellifica</i> | <i>A. bombi</i> | <i>N. ceranae</i>                  | –                       | <i>S. melliferum</i> |
| <i>H. truncorum</i> 1 | 5 July 2012   | BQCV                              | –                   | –               | <i>N. ceranae</i>                  | –                       | –                    |
| <i>H. truncorum</i> 2 | 5 July 2012   | –                                 | –                   | –               | <i>N. ceranae</i>                  | –                       | –                    |
| <i>H. truncorum</i> 3 | 5 July 2012   | BQCV                              | –                   | <i>A. bombi</i> | <i>N. ceranae</i>                  | –                       | –                    |

Hive 3, 5 and 10: hive identification number, ALPV: Aphid Lethal Paralysis Virus, *A. vaga*: *Andrena vaga*, *A. ventralis*: *Andrena ventralis*, BQCV: Black Queen Cell Virus, *C. bombi*: *Crithidia bombi*, *C. mellifica*: *Crithidia mellifica*, DWV: Deformed Wing Virus, *H. truncorum*: *Heriades truncorum*, LSV: Lake Sinai Virus, *N. apis*: *Nosema apis*, *N. ceranae*: *Nosema ceranae*, *N. thomsoni*: *Nosema thomsoni*, *O. bicornis*: *Osmia bicornis*, *O. cornuta*: *Osmia cornuta*, SBV: Sacbrood Virus, VdMLV: *Varroa destructor* Macula-like Virus.

(Qiagen). PCR products were electrophoresed using 1.4% agarose gels, stained with ethidium bromide and visualised under UV light.

Honey bee samples (1 µl RNA) were screened using the BeeDoctor tool, capable of detecting actin (honey bee control gene), Acute Bee Paralysis Virus (ABPV), BQCV, Chronic Bee Paralysis Virus (CBPV), DWV, SacBrood Virus (SBV) and Slow Bee Paralysis Virus (SBPV) (De Smet et al., 2012). ABPV probes amplify ABPV, Israeli Acute Paralysis Virus and Kashmir Bee Virus; DWV probes amplify DWV, Kakugo Virus and *Varroa destructor* Virus-1 (VDV-1). Solitary bees were analysed for these viruses using RT-PCR. Other viruses (Aphid Lethal Paralysis Virus strain Brookings (ALPV), AmFV, Big Sioux River Virus (BSRV), Lake Sinai Virus (LSV), Tobacco Ringspot Virus (TRSV) and *Varroa destructor* Macula-like Virus (VdMLV), bacteria (*Spiroplasma* spp.), fungi (*Ascosphaera* spp.) and *Nosema* spp.) and protists (*Apicystis bombi* and *Crithidia* spp.) were screened by PCR. The cytochrome c oxidase subunit I gene was used as control gene for the solitary bees.

#### 2.4. Sequence analysis

Amplicons were cloned using the TOPO TA Cloning Kit for Sequencing (Invitrogen). Plasmids and PCR products were sequenced using gene-specific or M13 primers. Sequences are deposited as AB859946–AB859948 (DWV), AB860145–AB860146 (*Crithidia* spp.), AB859949–AB859952 (VdMLV), HG764796–HG764797 (BQCV), HG764798–HG764799 (SBV), KF768348–KF68351 (LSV), KJ685944 (AmFV), KJ685945–KJ685947 (*Ascosphaera* spp.).


Phylogenetic trees were inferred via maximum likelihood (ML) using PhyML 3.0 (Guindon et al., 2010) with the Le-Gascuel (LG)

amino acid substitution model (Le and Gascuel, 2008) and approximate likelihood ratio test non-parametric branch support based on a Shimodaira-Hasegawa-like (aLRT SH-like) procedure (Anisimova and Gascuel, 2006).

### 3. Results and discussion

We were able to demonstrate the presence of AmFV in all samples. Most other viruses (BQCV, DWV, SBV and VdMLV) were detected both in honey bees and in a smaller subset of solitary bees (see Table 1). ALPV was detected in honey bee samples only, but the viruses ABPV, BSRV, CBPV, SBPV and TRSV were not discovered at all. Protists and fungi appeared to be pervasive, whereas spiroplasmas were found only scarcely.

The discovered AmFV sequences were mutually identical and highly similar to Baculovirus sequences from Swiss bees (Genbank: JF304814) and *V. destructor* mites (Cornman et al., 2010) (Fig. A.1). Further, amplicons of the ribonucleotide reductase small subunit and thymidylate synthase were identical to those found in *V. destructor* (Genbank: GU980896–GU980897). The LSV strains found in honey bees, *A. vaga*, *O. bicornis* and *O. cornuta* were almost identical (Genbank: KF768348–KF68350). However, a deviating LSV strain (Genbank: KF768351) was detected both in *A. ventralis* and in some honey bees. Their Orf1 sequences showed only 76.7% amino acid similarity with each other. DWV was detected in all honey bee hives and in *O. bicornis*. Their 5' untranslated region (UTR) was equal to published DWV sequences. Although the 5' UTR amplicons appeared mutually identical, phylogenetic analysis of the DWV L protein resulted in two divergent strains (Fig. 1). Strain e3 (Genbank: AB859948), only detected in honey bees,



**Fig. 1.** Phylogeny of DWV L protein amino acid sequences of various isolates, including VDV-1 and Kakugo Virus. The phylogenetic tree was constructed using the maximum likelihood method under the LG parameter. Each sequence is indicated by its Genbank accession number. Strain e3 (Genbank: AB859948) is designated in blue, strains e5 and e10 (Genbank: AB859946, AB859947) in green. The VDV-1 clade is showed in red. Branch support for each node is indicated by aLRT values.

was part of the DWV clade. The other strains e5 and e10 (Genbank: AB859946, AB859947), detected in (different) honey bees and in *O. bicornis*, appeared to lie between DWV and *V. destructor* Virus-1. The finding of VdMLV in several solitary bees was remarkable as they are not a known host of *V. destructor*, which is considered to be the primary vector (de Miranda et al., 2011). This indicates that transmission can occur by other means, like contaminated flowers (Singh et al., 2010).

Honey bee trypanosomatids were identified as *Crithidia mellificae* haplotype A (Morimoto et al., 2013) by their cytochrome *b* sequence (Genbank: AB860145). *O. bicornis* and *A. vaga* appeared to be infected with *Crithidia bombi* (Genbank: AB860146), hitherto only reported in bumble bees (Macfarlane et al., 1995). The neogregarine *A. bombi*, found in the present study in all bee species, was previously only discovered in social bees (Lipa and Triggiani, 1996). Sequencing of its internal transcribed spacer (ITS) region from isolates of honey bees and *O. cornuta* assigned them as haplotype Uni (Maharramov et al., 2013).

*Ascospaera apis* was detected in several honey bee samples. The other *Ascospaera* spp. that we discovered in *O. cornuta* and *H. truncorum* (Genbank: KJ685945–KJ685947) matched (~98%) *Ascospaera callicarpa* (Genbank: JX070046), previously reported from *Chelostoma florissomne* (Megachilidae) (Wynns et al., 2013). The related *Bettsia alvei* was also found in *O. cornuta*. Several sequences had no strong match (<93%) as reported before (Evison et al., 2012). The microsporidian parasite *N. ceranae* was detected in all species, except *A. vaga*. This gut parasite was previously also detected in bumble bees (Furst et al., 2014; Graystock et al., 2013; Li et al., 2012). Some honey bees were co-infected with *N. apis*.

Surprisingly, the microsporidium detected in *A. vaga* (Genbank: KC596023) appeared to be highly related (99.8%) to *N. thomsoni* and other microsporidia from the same clade (Li et al., 2012).

The spiroplasmas found in honey bees appeared to be *Spiroplasma melliferum*. *Spiroplasma apis* and *S. melliferum* were both discovered in *O. bicornis*, an unprecedented host (Clark et al., 1985; Mouches et al., 1983).

#### 4. Conclusions

Our study identified several honey bee pathogens in solitary bees living in the proximity of an apiary. Our results suggest that bee hives represent a putative source of pathogens for other pollinators. Similarly, solitary bees may act as a reservoir of honey bee pathogens.

#### Acknowledgment

This work was supported by the Research Foundation-Flanders (FWO, research Grant G.0628.11).

#### Appendix A. Supplementary material

Supplementary data associated with this article can be found, in the online version, at <http://dx.doi.org/10.1016/j.jip.2014.08.007>.

## References

- Anisimova, M., Gascuel, O., 2006. Approximate likelihood-ratio test for branches: a fast, accurate, and powerful alternative. *Syst. Biol.* 55, 539–552.
- Clark, T.B., Whitcomb, R.F., Tully, J.G., Mouches, C., Saillard, C., Bove, J.M., Wroblewski, H., Carle, P., Rose, D.L., Henegar, R.B., Williamson, D.L., 1985. *Spiroplasma melliferum*, a new species from the honeybee (*Apis mellifera*). *Int. J. Syst. Bacteriol.* 35, 296–308.
- Cornman, R.S., Schatz, M.C., Johnston, S.J., Chen, Y.P., Pettis, J., Hunt, G., Bourgeois, L., Elisk, C., Anderson, D., Grozinger, C.M., Evans, J.D., 2010. Genomic survey of the ectoparasitic mite *Varroa destructor*, a major pest of the honey bee *Apis mellifera*. *BMC Genom.* 11.
- De Smet, L., Ravoet, J., de Miranda, J.R., Wenseleers, T., Mueller, M.Y., Moritz, R.F.A., de Graaf, D.C., 2012. BeeDoctor, a versatile MLPA-based diagnostic tool for screening bee viruses. *PLoS One* 7, e47953.
- Evison, S.E., Roberts, K.E., Laurenson, L., Pietravalle, S., Hui, J., Biesmeijer, J.C., Smith, J.E., Budge, G., Hughes, W.O., 2012. Pervasiveness of parasites in pollinators. *PLoS One* 7, e30641.
- Fries, I., Paxton, R.J., Tengo, J., Slemenda, S.B., da Silva, A.J., Pieniazek, N.J., 1999. Morphological and molecular characterization of *Antonosporea scoticae* n. gen., n. sp. (Protozoa, Microsporidia) a parasite of the communal bee, *Andrena scotica* Perkins, 1916 (Hymenoptera, Andrenidae). *Eur. J. Protistol.* 35, 183–193.
- Furst, M.A., McMahon, D.P., Osborne, J.L., Paxton, R.J., Brown, M.J.F., 2014. Disease associations between honeybees and bumblebees as a threat to wild pollinators. *Nature* 506, 364–366.
- Graystock, P., Yates, K., Darvill, B., Goulson, D., Hughes, W.O., 2013. Emerging dangers: deadly effects of an emergent parasite in a new pollinator host. *J. Invertebr. Pathol.* 114, 114–119.
- Guindon, S., Dufayard, J.F., Lefort, V., Anisimova, M., Hordijk, W., Gascuel, O., 2010. New algorithms and methods to estimate maximum-likelihood phylogenies: assessing the performance of PhyML 3.0. *Syst. Biol.* 59, 307–321.
- Le, S.Q., Gascuel, O., 2008. An improved general amino acid replacement matrix. *Mol. Biol. Evol.* 25, 1307–1320.
- Levitt, A.L., Singh, R., Cox-Foster, D.L., Rajotte, E., Hoover, K., Ostiguy, N., Holmes, E.C., 2013. Cross-species transmission of honey bee viruses in associated arthropods. *Virus Res.* 176, 232–240.
- Li, J.L., Chen, W.F., Wu, J., Peng, W.J., An, J.D., Schmid-Hempel, P., Schmid-Hempel, R., 2012. Diversity of *Nosema* associated with bumblebees (*Bombus* spp.) from China. *Int. J. Parasitol.* 42, 49–61.
- Lipa, J.J., Triggiani, O., 1996. *Apicystis* gen. nov. and *Apicystis bombi* (Liu, Macfarlane & Pengelly) comb. nov. (Protozoa: Neogregarinida), a cosmopolitan parasite of *Bombus* and *Apis* (Hymenoptera: Apidae). *Apidologie* 27, 29–34.
- Macfarlane, R.P., Lipa, J.J., Liu, H.J., 1995. Bumble bee pathogens and internal enemies. *Bee World* 76, 130–148.
- Maharramov, J., Meeus, I., Maebe, K., Arbetman, M., Morales, C., Graystock, P., Hughes, W.O.H., Plischuk, S., Lange, C.E., de Graaf, D.C., Zapata, N., de la Rosa, J.J.P., Murray, T.E., Brown, M.J.F., Smagghe, G., 2013. Genetic variability of the neogregarine *Apicystis bombi*, an etiological agent of an emergent bumblebee disease. *PLoS One* 8, e81475.
- de Miranda, J.R., Gauthier, L., Ribière, M., Chen, Y., 2011. Honey bee viruses and their effect on bee and colony health. In: Sammataro, D., Yoder, J.A. (Eds.), *Honey Bee Colony Health: Challenges and Sustainable Solutions*. CRC Press, pp. 71–112.
- Morimoto, T., Kojima, Y., Yoshiyama, M., Kimura, K., Yang, B., Peng, G.D., Kadowaki, T., 2013. Molecular detection of protozoan parasites infecting *Apis mellifera* colonies in Japan. *Environ. Microbiol. Rep.* 5, 74–77.
- Mouches, C., Bove, J.M., Tully, J.G., Rose, D.L., McCoy, R.E., Carlejunca, P., Garnier, M., Saillard, C., 1983. *Spiroplasma apis*, a new species from the honeybee *Apis mellifera*. *Ann. Inst. Pasteur Mic.* A134, 383–397.
- Runckel, C., Flenniken, M.L., Engel, J.C., Ruby, J.G., Ganem, D., Andino, R., DeRisi, J.L., 2011. Temporal analysis of the honey bee microbiome reveals four novel viruses and seasonal prevalence of known viruses, *Nosema*, and *Crithidia*. *PLoS One* 6, e20656.
- Scheuchl, E., 1996. *Illustrierte Bestimmungstabellen der Wildbienen Deutschlands und Österreichs. Band II. Megachilidae – Melittidae*. Eigenverlag, Velden.
- Schmid-Egger, C., Scheuchl, E., 1997. *Illustrierte Bestimmungstabellen der Wildbienen Deutschlands und Österreichs unter Berücksichtigung der Arten der Schweiz. Band III. Andreninae*. Eigenverlag, Velden.
- Singh, R., Levitt, A.L., Rajotte, E.G., Holmes, E.C., Ostiguy, N., Vanengelsdorp, D., Lipkin, W.I., Depamphilis, C.W., Toth, A.L., Cox-Foster, D.L., 2010. RNA viruses in hymenopteran pollinators: evidence of inter-Taxa virus transmission via pollen and potential impact on non-*Apis* hymenopteran species. *PLoS One* 5, e14357.
- Westrich, P., 1990. *Die Wildbienen Baden-Württembergs. Spezieller Teil*. Eugen Ulmer Verlag, Stuttgart.
- Wynns, A.A., Jensen, A.B., Eilenberg, J., 2013. *Ascosphaera callicarpa*, a new species of bee-loving fungus, with a key to the genus for Europe. *PLoS One* 8, e73419.